

Climate Change Vulnerability of Trees Commonly Planted or Being Considered for Planting in the Twin Cities Region

DRAFT-November 14, 2016

Latin Name	Common Name	Planted Adapt Class	Natural Adapt Class	Effect from Zone Shift-low emissions	Effect from Zone Shift-high emissions	Tree Atlas-Low Emissions	Tree Atlas-High Emissions	Vulnerability-developed/planted sites	Vulnerability-natural areas
<i>Abies balsamea</i>	balsam fir	medium	na	negative	negative			moderate-high	
<i>Abies concolor</i>	white fir	medium	na	no effect	negative			moderate-high	
<i>Abies fraseri</i>	Fraser Fir	medium	na	no effect	negative			moderate-high	
<i>Abies koreana</i>	Korean Fir	medium	na	no effect	positive			low-moderate	
<i>Acer ginnala</i>	Amur maple	medium	na	no effect	negative			moderate-high	
<i>Acer negundo</i>	boxelder	medium	high			small decrease	small increase	moderate	low-moderate
<i>Acer nigrum</i>	black maple	medium	high			extirpated	extirpated	moderate-high	moderate
<i>Acer platanoides</i>	Norway maple	high	high	no effect	negative			moderate	moderate
<i>Acer rubrum</i>	red maple	medium	high			no change	small increase	low-moderate	low
<i>Acer saccharinum</i>	silver maple	medium	high			large increase	large increase	low-moderate	low
<i>Acer saccharum</i>	sugar maple	medium	high			no change	large decrease	moderate-high	moderate
<i>Acer tataricum</i>	tatarian maple	medium	na	no effect	negative			moderate-high	
<i>Acer triflorum</i>	threeflower maple	medium	na	positive	negative			moderate	
<i>Acer truncatum</i>	Shantung maple	high	na	positive	positive			low	
<i>Acer x freemanii</i>	Freeman maple	high	na	no effect	no effect			low	
<i>Aescleus glabra</i>	Ohio buckeye	medium	low			NA	new habitat	low-moderate	moderate
<i>Aescleus hippocastanatum</i>	horsechestnut	medium	na	no effect	no effect			low-moderate	
<i>Aesculus flava (octandra)</i>	yellow buckeye	medium	na	no effect	no effect			low-moderate	
<i>Aesculus pavia</i>	red buckeye	high	na	positive	positive			low	
<i>Amelanchier arborea</i>	downy serviceberry	high	high	no effect	no effect			low	low
<i>Amelanchier laevis</i>	Allegheny serviceberry	high	high	no effect	no effect			low	low
<i>Betula alleghaniensis</i>	yellow birch	high	high			extirpated	extirpated	moderate	moderate
<i>Betula nigra</i>	river birch	medium	medium			small increase	small decrease	moderate	moderate
<i>Betula papyrifera</i>	paper birch	medium	medium			large decrease	large decrease	moderate-high	moderate-high
<i>Betula pendula</i>	European White birch	low	na	no effect	negative			moderate-high	
<i>Betula populifolia</i>	gray birch	low	medium	negative	negative			high	
<i>Carpinus caroliniana</i>	American hornbeam	high	high			no change	small increase	low	low
<i>Carya cordiformis</i>	bitternut hickory	medium	medium			large increase	small increase	low-moderate	low-moderate
<i>Carya glabra</i>	pignut hickory	medium	high			new habitat	new habitat	low-moderate	low
<i>Carya illinoensis</i>	pecan	low	na			NA	new habitat	moderate	
<i>Carya laciniosa</i>	shellbark hickory	low	medium	positive	positive			moderate	
<i>Carya ovata</i>	shagbark hickory	medium	high			new habitat	new habitat	low-moderate	

*Vulnerability for ash species is considered high due to emerald ash borer

Tree Atlas model projections from: <http://www.treesearch.fs.fed.us/pubs/13412>

Adaptation scores based on: <http://treesearch.fs.fed.us/pubs/38643>

Climate Change Vulnerability of Trees Commonly Planted or Being Considered for Planting in the Twin Cities Region
DRAFT-November 14, 2016

Latin Name	Common Name	Planted Adapt Class	Natural Adapt Class	Effect from Zone Shift-low emissions	Effect from Zone Shift-high emissions	Tree Atlas-Low Emissions	Tree Atlas-High Emissions	Vulnerability-developed/planted sites	Vulnerability-natural areas
<i>Catalpa ovata</i>	Chinese catalpa	medium	na	positive	positive			low-moderate	
<i>Catalpa speciosa</i>	northern catalpa	low	low	no effect	no effect			moderate	
<i>Celtis laevigata</i>	sugarberry	medium	na			NA	new habitat	low-moderate	
<i>Celtis occidentalis</i>	northern hackberry	high	high			large increase	large increase	low	low
<i>Cercidiphyllum japonicum</i>	Katsura tree	low	na	no effect	no effect			moderate	
<i>Cercis canadensis</i>	eastern redbud	medium	high			NA	new habitat	low-moderate	low
<i>Cladrastis kentukea</i>	yellowwood	high	na	no effect	no effect			low	
<i>Cornus alternifolia</i>	pagoda dogwood	medium	medium	no effect	negative			moderate	moderate
<i>Crataegus acutifolia</i>	cockspur hawthorn	high	na	no effect	negative			moderate	
<i>Fagus grandifolia</i>	American beech	medium	medium			new habitat	new habitat	low-moderate	
<i>Fagus sylvatica</i>	European beech	medium	na	no effect	no effect			low-moderate	low-moderate
<i>Fraxinus americana</i>	white ash	low	medium			small increase	large increase	high*	high*
<i>Fraxinus nigra</i>	black Ash	low	low			small decrease	small decrease	high*	high*
<i>Fraxinus pennsylvanica</i>	Green ash	medium	medium			small decrease	no change	high*	high*
<i>Ginkgo biloba</i>	gingko/maidenhair tree	high	na	no effect	no effect			low	low
<i>Gleditsia triacanthos</i>	honeylocust	medium	high			small increase	small increase	low-moderate	low
<i>Gymnocladus dioicus</i>	Kentucky coffeetree	high	na	no effect	no effect			low	
<i>Juglans nigra</i>	black walnut	medium	medium			large increase	large increase	low-moderate	low-moderate
<i>Juniperus virginiana</i>	eastern redcedar	high	medium			large increase	large increase	low	low-moderate
<i>Liriodendron tulipifera</i>	tuliptree	low	na	positive	positive			moderate	
<i>Maackia amurensis</i>	Amur maackia	high	na	no effect	negative			moderate	
<i>Maclura pomifera</i>	Osage orange	high	na			new habitat	new habitat	low	
<i>Magnolia acuminata</i>	cucumbertree	medium	na	positive	positive			low-moderate	
<i>Magnolia stellata</i>	star magnolia	medium	na	no effect	no effect			moderate	
<i>Malus spp</i>	crabapple species	medium	na	no effect	no effect			moderate	
<i>Metasequoia glyptostroboides</i>	dawn redwood	medium	na	positive	positive			low-moderate	
<i>Morus alba</i>	white mulberry	medium	high	no effect	no effect			moderate	
<i>Morus rubra</i>	red mulberry	medium	medium			new habitat	new habitat	low-moderate	
<i>Ostrya virginiana</i>	Ironwood (easternhophornbe	high	high			no change	no change	low	low
<i>Phellodendron amurense</i>	Amur corktree	medium	na	no effect	no effect			moderate	
<i>Picea abies</i>	Norway spruce	medium	na	no effect	no effect			moderate	

*Vulnerability for ash species is considered high due to emerald ash borer

Tree Atlas model projections from: <http://www.treesearch.fs.fed.us/pubs/13412>

Adaptation scores based on: <http://treesearch.fs.fed.us/pubs/38643>

Climate Change Vulnerability of Trees Commonly Planted or Being Considered for Planting in the Twin Cities Region

DRAFT-November 14, 2016

Latin Name	Common Name	Planted Adapt Class	Natural Adapt Class	Effect from Zone Shift-low emissions	Effect from Zone Shift-high emissions	Tree Atlas-Low Emissions	Tree Atlas-High Emissions	Vulnerability-developed/planted sites	Vulnerability-natural areas
<i>Picea canadensis (glauca)</i>	White spruce	medium	na			no change	no change	moderate	
<i>Picea glauca var. densata</i>	Black Hills spruce	medium	na	no effect	negative			moderate-high	
<i>Picea pungens</i>	blue spruce	medium	na	no effect	no effect			moderate	
<i>Pinus banksiana</i>	jack pine	low	na			small decrease	small decrease	high	
<i>Pinus nigra</i>	Austrian pine	medium	na	no effect	no effect			moderate	
<i>Pinus ponderosa</i>	ponderosa Pine	low	na	no effect	no effect			moderate-high	
<i>Pinus resinosa</i>	red pine	low	na			small increase	no change	moderate	
<i>Pinus strobus</i>	white pine	low	medium			no change	large decrease	moderate-high	
<i>Pinus sylvestris</i>	Scotch pine	medium	na	no effect	negative			moderate-high	
<i>Platanus occidentalis</i>	American sycamore	medium	na			NA	new habitat	low-moderate	
<i>Platanus x acerifolia</i>	planetree	medium	na	no effect	no effect			low-moderate	low-moderate
<i>Populus aurea (tremuloides)</i>	quaking aspen	low	medium			small decrease	large decrease	high	moderate-high
<i>Populus balsamifera</i>	balsam poplar	low	medium			no change	small increase	moderate	low-moderate
<i>Populus deltoides</i>	cottonwood	low	medium			small increase	large increase	moderate	low-moderate
<i>Populus grandidentata</i>	big-toothed aspen	low	high			small decrease	large decrease	high	moderate
<i>Prunus americana</i>	American plum	medium	medium			small increase	large increase	low-moderate	low-moderate
<i>Prunus cerasus</i>	sour Cherry	medium	na	no effect	no effect			moderate	
<i>Prunus maackii</i>	Amur cherry	high	na	no effect	no effect			low	
<i>Prunus mandshurica</i>	apricot (Manchurian)	low	na	no effect	no effect			moderate-high	
<i>Prunus pensylvanica</i>	pin cherry	low	medium			extirpated	extirpated	high	moderate-high
<i>Prunus serotina</i>	black cherry	low	medium			small increase	small decrease	high	moderate-high
<i>Prunus serrulata 'Kwanzan'</i>	Kwanzan Cherry	low	na	positive	positive			moderate	
<i>Prunus virginiana</i>	common chokecherry	medium	medium			small decrease	extirpated	moderate-high	moderate-high
<i>Prunus x cistena</i>	purpleleaf Sand Cherry	medium	na	no effect	no effect			moderate	
<i>Pseudotsuga mucronata</i>	douglas fir	low	na	no effect	negative			high	
<i>Pyrus spp.</i>	pear	medium	na	no effect	no effect			moderate	
<i>Pyrus ussuriensis</i>	Ussurian Pear	low	na	no effect	no effect			moderate-high	
<i>Quercus alba</i>	white oak	medium	high			small increase	no change	low-moderate	low
<i>Quercus bicolor</i>	swamp white oak	high	medium			no change	small increase	low	low-moderate
<i>Quercus coccinea</i>	scarlet oak	high	na	positive	positive			low	
<i>Quercus ellipsoidalis</i>	northern pin oak (Hill's Oak)	medium	medium			no change	small decrease	moderate	moderate

*Vulnerability for ash species is considered high due to emerald ash borer

Tree Atlas model projections from: <http://www.treesearch.fs.fed.us/pubs/13412>

Adaptation scores based on: <http://treesearch.fs.fed.us/pubs/38643>

Climate Change Vulnerability of Trees Commonly Planted or Being Considered for Planting in the Twin Cities Region

DRAFT-November 14, 2016

Latin Name	Common Name	Planted Adapt Class	Natural Adapt Class	Effect from Zone Shift-low emissions	Effect from Zone Shift-high emissions	Tree Atlas-Low Emissions	Tree Atlas-High Emissions	Vulnerability-developed/planted sites	Vulnerability-natural areas
<i>Quercus imbricaria</i>	shingle oak	high	na			NA	new habitat	low	
<i>Quercus macrocarpa</i>	bur oak	high	high			no change	no change	low	low
<i>Quercus muehlenbergii</i>	chinkapin oak	medium	na			new habitat	new habitat	low-moderate	
<i>Quercus palustris</i>	pin oak	medium	medium			NA	new habitat	low-moderate	low-moderate
<i>Quercus prinus</i>	chestnut oak	high	high	positive	positive			low	low
<i>Quercus rubra</i>	northern red oak	high	high			small decrease	large decrease	moderate	moderate
<i>Quercus velutina</i>	black oak	medium	high			large increase	small increase	low-moderate	low
<i>Quercus x macdanielii</i>	heritage oak	high	na	no effect	no effect			low	
<i>Rhamnus cathartica</i>	European buckthorn	high	high	no effect	no effect			low	low
<i>Rhus typhina</i>	staghorn sumac	medium	high	no effect	no effect			moderate	low
<i>Robinia pseudoacacia</i>	black locust	medium	high			large increase	large increase	low-moderate	low
<i>Salix amygdaloides</i>	peachleaf willow	low	low			no change	small increase	moderate-high	moderate-high
<i>Salix babylonica</i>	weeping willow	medium	na	positive	positive			low-moderate	
<i>Salix discolor</i>	pussy willow	low	medium	no effect	no effect			moderate-high	moderate-high
<i>Salix nigra</i>	black willow	low	Low			no change	small increase	moderate	moderate
<i>Sorbus alnifolia</i>	Korean mountain-ash	medium	na	no effect	no effect			moderate	
<i>Sorbus aucoparia</i>	European Mountain Ash	medium	na	no effect	negative			moderate-high	
<i>Sorbus decora</i>	showy Mountain Ash	low	na	negative	negative			high	
<i>Syringa reticulata</i>	Japanese tree lilac	high	na	no effect	negative			moderate	
<i>Thuja occidentalis</i>	northern white cedar	medium	medium			no change	small increase	low-moderate	
<i>Tilia americana</i>	American basswood	medium	medium			small decrease	small decrease	moderate-high	moderate-high
<i>Tilia cordata</i>	little leaf linden	high	na	no effect	no effect			low	
<i>Tilia x euchlora</i>	Crimean linden	high	na	no effect	negative			moderate	
<i>Ulmus americana</i>	American elm	medium	high			small increase	small increase	low-moderate	low
<i>Ulmus spp</i>	elm cultivars	high	na	no effect	no effect			low	
<i>Ulmus pumila</i>	Siberian elm	medium	high	no effect	no effect			low-moderate	low
<i>Ulmus rubra</i>	slippery elm	medium	medium			no change	small increase	low-moderate	low-moderate
<i>Viburnum lentago</i>	nannyberry	high	high	no effect	no effect			low	low

*Vulnerability for ash species is considered high due to emerald ash borer

Tree Atlas model projections from: <http://www.treesearch.fs.fed.us/pubs/13412>

Adaptation scores based on: <http://treesearch.fs.fed.us/pubs/38643>